



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 04-08-2023

Weather Forecast of **ANDANALLUR** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2023-08-04(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-05	2023-08-06	2023-08-07	2023-08-08	2023-08-09
Rainfall	0.0	0.0	0.0	3.0	2.0
Tmax(°C)	33.0	33.0	32.0	33.0	34.0
Tmin(°C)	23.0	23.0	23.0	23.0	23.0
RH-I(%)	61	60	62	60	59
RH-II(%)	51	48	47	46	46
Wind Speed(kmph)	11	11	11	7	9
Wind Direction(Degree)	255	278	287	282	283
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Rainfall is expected for 2 day in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (60.0% & 48.0%), 33.0°C and 23.0°C respectively. The clear sky may appear. Expected average wind speed is 9.8 kmph. The Wind is expected to flow in west direction.

General Advisory:

Tamilnadu's average rainfall is likely to be increase than the normal during this week

SMS Advisory:

Blackgram VBN 8 & VBN 11 are available in KVK, Sirugamani, Trichy – 115. Contact: 8508835287

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing weather thrips infestation may occur in paddy at nursery stage. To control thrips infestation spray thiamethoxam 25% WG 4g
SUGARCANE	To control mealy bug in sugarcane, spray malathion 2ml/litre.
COTTON	Due to prevailing weather the white fly infestation may occur in cotton. To control white fly in cotton, Spray profenophos 50% EC 1000ml/ha.
MAIZE	Due to prevailing weather the downy mildew incidence may occur in maize. To control downy mildew, Spray mancozeb 1 kg/ha at 20 days after sowing.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Due to prevailing weather the white fly incidence may occur in Tomato. To control install yellow sticky traps @12/ha.
BANANA	Light to heavy wind/gust speed will expected in forthcoming days so farmers are advised to save the banana crop by using stacking method with casuarina poles.

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	Due to prevailing weather the Thrips incidence may occur in Chillies. To control, Spray Imidacloprid 17.8 SL 3 ml/10 litre.
JASMINE	For red spider mite incidence in jasmine, spraying of fenazagnin 10%EC @ 2ml/litre is recommended.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Pneumonia in goats – Fever, Respiratory distress, Mucous discharge from nostril, Reduced feed intake & Weight gain and Cough. Remedial measures: Provide clean water and ventilated house.
COW	To avoid mastitis disease in milching animals use potassium permanganate for cleaning the udder before and after milking.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Due to high wind speed feed material of poultries may be drifted hence hanging of gunny bags at the borders of the shed to reduce the wind speed.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 04-08-2023

Weather Forecast of **LALGUDI** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2023-08-04(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-05	2023-08-06	2023-08-07	2023-08-08	2023-08-09
Rainfall	0.0	0.0	0.0	2.0	0.0
Tmax(°C)	37.0	36.0	34.0	35.0	37.0
Tmin(°C)	25.0	25.0	25.0	25.0	25.0
RH-I(%)	60	59	61	62	61
RH-II(%)	44	40	43	42	39
Wind Speed(kmph)	16	15	14	10	10
Wind Direction(Degree)	262	278	282	267	290
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Rainfall is expected for 1 day in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (61.0% & 42.0%), 36.0°C and 25.0°C respectively. The clear sky may appear. Expected average wind speed is 13 kmph. The Wind is expected to flow in west direction.

General Advisory:

Tamilnadu's average rainfall is likely to be increase than the normal during this week

SMS Advisory:

Blackgram VBN 8 & VBN 11 are available in KVK, Sirugamani, Trichy – 115. Contact: 8508835287

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing weather thrips infestation may occur in paddy at nursery stage. To control thrips infestation spray thiamethoxam 25% WG 4g
SUGARCANE	To control mealy bug in sugarcane, spray malathion 2ml/litre.
COTTON	Due to prevailing weather the white fly infestation may occur in cotton. To control white fly in cotton, Spray profenophos 50% EC 1000ml/ha.
MAIZE	Due to prevailing weather the downy mildew incidence may occur in maize. To control downy mildew, Spray mancozeb 1 kg/ha at 20 days after sowing.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Due to prevailing weather the white fly incidence may occur in Tomato. To control install yellow sticky traps @12/ha.
BANANA	Light to heavy wind/gust speed will expected in forthcoming days so farmers are advised to save the banana crop by using stacking method with casuarina poles.

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	Due to prevailing weather the Thrips incidence may occur in Chillies. To control, Spray Imidacloprid 17.8 SL 3 ml/10 litre.
JASMINE	For red spider mite incidence in jasmine, spraying of fenazagnin 10%EC @ 2ml/litre is recommended.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Pneumonia in goats – Fever, Respiratory distress, Mucous discharge from nostril, Reduced feed intake & Weight gain and Cough. Remedial measures: Provide clean water and ventilated house.
COW	To avoid mastitis disease in milching animals use potassium permanganate for cleaning the udder before and after milking.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Due to high wind speed feed material of poultries may be drifted hence hanging of gunny bags at the borders of the shed to reduce the wind speed.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 04-08-2023

Weather Forecast of **MANACHANELLUR** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2023-08-04(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-05	2023-08-06	2023-08-07	2023-08-08	2023-08-09
Rainfall	0.0	0.0	0.0	2.0	0.0
Tmax(°C)	37.0	36.0	34.0	35.0	37.0
Tmin(°C)	25.0	25.0	25.0	25.0	25.0
RH-I(%)	60	59	61	62	61
RH-II(%)	44	40	43	41	39
Wind Speed(kmph)	16	15	14	10	10
Wind Direction(Degree)	263	279	282	267	289
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Rainfall is expected for 1 day in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (61.0% & 41.0%), 36.0°C and 25.0°C respectively. The clear sky may appear. Expected average wind speed is 13 kmph. The Wind is expected to flow in west direction.

General Advisory:

Tamilnadu's average rainfall is likely to be increase than the normal during this week

SMS Advisory:

Blackgram VBN 8 & VBN 11 are available in KVK, Sirugamani, Trichy – 115. Contact: 8508835287

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing weather thrips infestation may occur in paddy at nursery stage. To control thrips infestation spray thiamethoxam 25% WG 4g
SUGARCANE	To control mealy bug in sugarcane, spray malathion 2ml/litre.
COTTON	Due to prevailing weather the white fly infestation may occur in cotton. To control white fly in cotton, Spray profenophos 50% EC 1000ml/ha.
MAIZE	Due to prevailing weather the downy mildew incidence may occur in maize. To control downy mildew, Spray mancozeb 1 kg/ha at 20 days after sowing.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Due to prevailing weather the white fly incidence may occur in Tomato. To control install yellow sticky traps @12/ha.
BANANA	Light to heavy wind/gust speed will expected in forthcoming days so farmers are advised to save the banana crop by using stacking method with casuarina poles.

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	Due to prevailing weather the Thrips incidence may occur in Chillies. To control, Spray Imidacloprid 17.8 SL 3 ml/10 litre.
JASMINE	For red spider mite incidence in jasmine, spraying of fenazagnin 10%EC @ 2ml/litre is recommended.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Pneumonia in goats – Fever, Respiratory distress, Mucous discharge from nostril, Reduced feed intake & Weight gain and Cough. Remedial measures: Provide clean water and ventilated house.
COW	To avoid mastitis disease in milching animals use potassium permanganate for cleaning the udder before and after milking.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Due to high wind speed feed material of poultries may be drifted hence hanging of gunny bags at the borders of the shed to reduce the wind speed.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 04-08-2023

Weather Forecast of **MANAPPARAI** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2023-08-04(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-05	2023-08-06	2023-08-07	2023-08-08	2023-08-09
Rainfall	0.0	0.0	0.0	2.0	1.0
Tmax(°C)	36.0	35.0	34.0	35.0	36.0
Tmin(°C)	25.0	24.0	25.0	24.0	24.0
RH-I(%)	59	59	60	61	59
RH-II(%)	47	43	44	44	42
Wind Speed(kmph)	15	14	13	9	10
Wind Direction(Degree)	258	278	286	278	291
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Rainfall is expected for 2 day in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (60.0% & 44.0%), 35.0°C and 24.0°C respectively. The clear sky may appear. Expected average wind speed is 12.2 kmph. The Wind is expected to flow in west direction.

General Advisory:

Tamilnadu's average rainfall is likely to be increase than the normal during this week

SMS Advisory:

Blackgram VBN 8 & VBN 11 are available in KVK, Sirugamani, Trichy – 115. Contact: 8508835287

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing weather thrips infestation may occur in paddy at nursery stage. To control thrips infestation spray thiamethoxam 25% WG 4g
SUGARCANE	To control mealy bug in sugarcane, spray malathion 2ml/litre.
COTTON	Due to prevailing weather the white fly infestation may occur in cotton. To control white fly in cotton, Spray profenophos 50% EC 1000ml/ha.
MAIZE	Due to prevailing weather the downy mildew incidence may occur in maize. To control downy mildew, Spray mancozeb 1 kg/ha at 20 days after sowing.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Due to prevailing weather the white fly incidence may occur in Tomato. To control install yellow sticky traps @12/ha.
BANANA	Light to heavy wind/gust speed will expected in forthcoming days so farmers are advised to save the banana crop by using stacking method with casuarina poles.

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	Due to prevailing weather the Thrips incidence may occur in Chillies. To control, Spray Imidacloprid 17.8 SL 3 ml/10 litre.
JASMINE	For red spider mite incidence in jasmine, spraying of fenazagnin 10%EC @ 2ml/litre is recommended.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Pneumonia in goats – Fever, Respiratory distress, Mucous discharge from nostril, Reduced feed intake & Weight gain and Cough. Remedial measures: Provide clean water and ventilated house.
COW	To avoid mastitis disease in milching animals use potassium permanganate for cleaning the udder before and after milking.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Due to high wind speed feed material of poultries may be drifted hence hanging of gunny bags at the borders of the shed to reduce the wind speed.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 04-08-2023

Weather Forecast of **MANIKANDAM** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2023-08-04(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-05	2023-08-06	2023-08-07	2023-08-08	2023-08-09
Rainfall	0.0	0.0	0.0	1.0	0.0
Tmax(°C)	37.0	36.0	34.0	36.0	38.0
Tmin(°C)	25.0	25.0	25.0	25.0	25.0
RH-I(%)	60	59	61	62	61
RH-II(%)	43	39	43	40	38
Wind Speed(kmph)	17	16	15	10	11
Wind Direction(Degree)	265	280	283	268	291
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Rainfall is expected for 1 day in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (61.0% & 41.0%), 36.0°C and 25.0°C respectively. The clear sky may appear. Expected average wind speed is 13.8 kmph. The Wind is expected to flow in west direction.

General Advisory:

Tamilnadu's average rainfall is likely to be increase than the normal during this week

SMS Advisory:

Blackgram VBN 8 & VBN 11 are available in KVK, Sirugamani, Trichy – 115. Contact: 8508835287

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing weather thrips infestation may occur in paddy at nursery stage. To control thrips infestation spray thiamethoxam 25% WG 4g
SUGARCANE	To control mealy bug in sugarcane, spray malathion 2ml/litre.
COTTON	Due to prevailing weather the white fly infestation may occur in cotton. To control white fly in cotton, Spray profenophos 50% EC 1000ml/ha.
MAIZE	Due to prevailing weather the downy mildew incidence may occur in maize. To control downy mildew, Spray mancozeb 1 kg/ha at 20 days after sowing.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Due to prevailing weather the white fly incidence may occur in Tomato. To control install yellow sticky traps @12/ha.
BANANA	Light to heavy wind/gust speed will expected in forthcoming days so farmers are advised to save the banana crop by using stacking method with casuarina poles.

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	Due to prevailing weather the Thrips incidence may occur in Chillies. To control, Spray Imidacloprid 17.8 SL 3 ml/10 litre.
JASMINE	For red spider mite incidence in jasmine, spraying of fenazagnin 10%EC @ 2ml/litre is recommended.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Pneumonia in goats – Fever, Respiratory distress, Mucous discharge from nostril, Reduced feed intake & Weight gain and Cough. Remedial measures: Provide clean water and ventilated house.
COW	To avoid mastitis disease in milching animals use potassium permanganate for cleaning the udder before and after milking.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Due to high wind speed feed material of poultries may be drifted hence hanging of gunny bags at the borders of the shed to reduce the wind speed.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 04-08-2023

Weather Forecast of **MARUNGAPURI** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2023-08-04(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-05	2023-08-06	2023-08-07	2023-08-08	2023-08-09
Rainfall	0.0	0.0	0.0	3.0	1.0
Tmax(°C)	36.0	36.0	34.0	36.0	37.0
Tmin(°C)	24.0	24.0	24.0	24.0	24.0
RH-I(%)	61	60	63	64	62
RH-II(%)	48	43	44	44	43
Wind Speed(kmph)	16	15	15	11	10
Wind Direction(Degree)	260	274	277	266	282
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Rainfall is expected for 2 day in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (62.0% & 44.0%), 36.0°C and 24.0°C respectively. The clear sky may appear. Expected average wind speed is 13.6 kmph. The Wind is expected to flow in west direction.

General Advisory:

Tamilnadu's average rainfall is likely to be increase than the normal during this week

SMS Advisory:

Blackgram VBN 8 & VBN 11 are available in KVK, Sirugamani, Trichy – 115. Contact: 8508835287

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing weather thrips infestation may occur in paddy at nursery stage. To control thrips infestation spray thiamethoxam 25% WG 4g
SUGARCANE	To control mealy bug in sugarcane, spray malathion 2ml/litre.
COTTON	Due to prevailing weather the white fly infestation may occur in cotton. To control white fly in cotton, Spray profenophos 50% EC 1000ml/ha.
MAIZE	Due to prevailing weather the downy mildew incidence may occur in maize. To control downy mildew, Spray mancozeb 1 kg/ha at 20 days after sowing.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Due to prevailing weather the white fly incidence may occur in Tomato. To control install yellow sticky traps @12/ha.
BANANA	Light to heavy wind/gust speed will expected in forthcoming days so farmers are advised to save the banana crop by using stacking method with casuarina poles.

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	Due to prevailing weather the Thrips incidence may occur in Chillies. To control, Spray Imidacloprid 17.8 SL 3 ml/10 litre.
JASMINE	For red spider mite incidence in jasmine, spraying of fenazagnin 10%EC @ 2ml/litre is recommended.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Pneumonia in goats – Fever, Respiratory distress, Mucous discharge from nostril, Reduced feed intake & Weight gain and Cough. Remedial measures: Provide clean water and ventilated house.
COW	To avoid mastitis disease in milching animals use potassium permanganate for cleaning the udder before and after milking.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Due to high wind speed feed material of poultries may be drifted hence hanging of gunny bags at the borders of the shed to reduce the wind speed.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 04-08-2023

Weather Forecast of **MUSIRI** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2023-08-04(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-05	2023-08-06	2023-08-07	2023-08-08	2023-08-09
Rainfall	0.0	0.0	0.0	3.0	1.0
Tmax(°C)	36.0	36.0	34.0	35.0	37.0
Tmin(°C)	24.0	24.0	25.0	24.0	24.0
RH-I(%)	61	60	63	63	62
RH-II(%)	47	42	44	44	42
Wind Speed(kmph)	16	15	14	11	10
Wind Direction(Degree)	260	275	278	266	284
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Rainfall is expected for 2 day in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (62.0% & 44.0%), 36.0°C and 24.0°C respectively. The clear sky may appear. Expected average wind speed is 13.2 kmph. The Wind is expected to flow in west direction.

General Advisory:

Tamilnadu's average rainfall is likely to be increase than the normal during this week

SMS Advisory:

Blackgram VBN 8 & VBN 11 are available in KVK, Sirugamani, Trichy – 115. Contact: 8508835287

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing weather thrips infestation may occur in paddy at nursery stage. To control thrips infestation spray thiamethoxam 25% WG 4g
SUGARCANE	To control mealy bug in sugarcane, spray malathion 2ml/litre.
COTTON	Due to prevailing weather the white fly infestation may occur in cotton. To control white fly in cotton, Spray profenophos 50% EC 1000ml/ha.
MAIZE	Due to prevailing weather the downy mildew incidence may occur in maize. To control downy mildew, Spray mancozeb 1 kg/ha at 20 days after sowing.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Due to prevailing weather the white fly incidence may occur in Tomato. To control install yellow sticky traps @12/ha.
BANANA	Light to heavy wind/gust speed will expected in forthcoming days so farmers are advised to save the banana crop by using stacking method with casuarina poles.

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	Due to prevailing weather the Thrips incidence may occur in Chillies. To control, Spray Imidacloprid 17.8 SL 3 ml/10 litre.
JASMINE	For red spider mite incidence in jasmine, spraying of fenazagnin 10%EC @ 2ml/litre is recommended.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Pneumonia in goats – Fever, Respiratory distress, Mucous discharge from nostril, Reduced feed intake & Weight gain and Cough. Remedial measures: Provide clean water and ventilated house.
COW	To avoid mastitis disease in milching animals use potassium permanganate for cleaning the udder before and after milking.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Due to high wind speed feed material of poultries may be drifted hence hanging of gunny bags at the borders of the shed to reduce the wind speed.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 04-08-2023

Weather Forecast of **PULLAMBADI** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2023-08-04(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-05	2023-08-06	2023-08-07	2023-08-08	2023-08-09
Rainfall	0.0	0.0	0.0	2.0	0.0
Tmax(°C)	36.0	36.0	34.0	35.0	37.0
Tmin(°C)	25.0	25.0	25.0	25.0	25.0
RH-I(%)	61	59	61	63	62
RH-II(%)	44	41	43	43	40
Wind Speed(kmph)	16	14	14	10	10
Wind Direction(Degree)	261	277	281	265	288
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Rainfall is expected for 1 day in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (61.0% & 41.0%), 36.0°C and 25.0°C respectively. The clear sky may appear. Expected average wind speed is 12.8 kmph. The Wind is expected to flow in west direction.

General Advisory:

Tamilnadu's average rainfall is likely to be increase than the normal during this week

SMS Advisory:

Blackgram VBN 8 & VBN 11 are available in KVK, Sirugamani, Trichy – 115. Contact: 8508835287

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing weather thrips infestation may occur in paddy at nursery stage. To control thrips infestation spray thiamethoxam 25% WG 4g
SUGARCANE	To control mealy bug in sugarcane, spray malathion 2ml/litre.
COTTON	Due to prevailing weather the white fly infestation may occur in cotton. To control white fly in cotton, Spray profenophos 50% EC 1000ml/ha.
MAIZE	Due to prevailing weather the downy mildew incidence may occur in maize. To control downy mildew, Spray mancozeb 1 kg/ha at 20 days after sowing.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Due to prevailing weather the white fly incidence may occur in Tomato. To control install yellow sticky traps @12/ha.
BANANA	Light to heavy wind/gust speed will expected in forthcoming days so farmers are advised to save the banana crop by using stacking method with casuarina poles.

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	Due to prevailing weather the Thrips incidence may occur in Chillies. To control, Spray Imidacloprid 17.8 SL 3 ml/10 litre.
JASMINE	For red spider mite incidence in jasmine, spraying of fenazagnin 10%EC @ 2ml/litre is recommended.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Pneumonia in goats – Fever, Respiratory distress, Mucous discharge from nostril, Reduced feed intake & Weight gain and Cough. Remedial measures: Provide clean water and ventilated house.
COW	To avoid mastitis disease in milching animals use potassium permanganate for cleaning the udder before and after milking.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Due to high wind speed feed material of poultries may be drifted hence hanging of gunny bags at the borders of the shed to reduce the wind speed.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 04-08-2023

Weather Forecast of **TATTAYANGARPETTAI** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2023-08-04(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-05	2023-08-06	2023-08-07	2023-08-08	2023-08-09
Rainfall	0.0	1.0	0.0	4.0	2.0
Tmax(°C)	36.0	36.0	34.0	35.0	37.0
Tmin(°C)	24.0	24.0	25.0	24.0	24.0
RH-I(%)	64	62	65	66	65
RH-II(%)	49	43	45	46	44
Wind Speed(kmph)	15	13	13	10	9
Wind Direction(Degree)	259	272	274	261	280
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Rainfall is expected for 2 day in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (64.0% & 45.0%), 36.0°C and 24.0°C respectively. The clear sky may appear. Expected average wind speed is 12 kmph. The Wind is expected to flow in west direction.

General Advisory:

Tamilnadu's average rainfall is likely to be increase than the normal during this week

SMS Advisory:

Blackgram VBN 8 & VBN 11 are available in KVK, Sirugamani, Trichy – 115. Contact: 8508835287

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing weather thrips infestation may occur in paddy at nursery stage. To control thrips infestation spray thiamethoxam 25% WG 4g
SUGARCANE	To control mealy bug in sugarcane, spray malathion 2ml/litre.
COTTON	Due to prevailing weather the white fly infestation may occur in cotton. To control white fly in cotton, Spray profenophos 50% EC 1000ml/ha.
MAIZE	Due to prevailing weather the downy mildew incidence may occur in maize. To control downy mildew, Spray mancozeb 1 kg/ha at 20 days after sowing.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Due to prevailing weather the white fly incidence may occur in Tomato. To control install yellow sticky traps @12/ha.
BANANA	Light to heavy wind/gust speed will expected in forthcoming days so farmers are advised to save the banana crop by using stacking method with casuarina poles.

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	Due to prevailing weather the Thrips incidence may occur in Chillies. To control, Spray Imidacloprid 17.8 SL 3 ml/10 litre.
JASMINE	For red spider mite incidence in jasmine, spraying of fenazagnin 10%EC @ 2ml/litre is recommended.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Pneumonia in goats – Fever, Respiratory distress, Mucous discharge from nostril, Reduced feed intake & Weight gain and Cough. Remedial measures: Provide clean water and ventilated house.
COW	To avoid mastitis disease in milching animals use potassium permanganate for cleaning the udder before and after milking.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Due to high wind speed feed material of poultries may be drifted hence hanging of gunny bags at the borders of the shed to reduce the wind speed.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 04-08-2023

Weather Forecast of **THIRUVERAMBUR** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2023-08-04(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-05	2023-08-06	2023-08-07	2023-08-08	2023-08-09
Rainfall	0.0	0.0	0.0	2.0	0.0
Tmax(°C)	37.0	36.0	34.0	35.0	37.0
Tmin(°C)	25.0	25.0	25.0	25.0	25.0
RH-I(%)	60	59	61	62	61
RH-II(%)	43	40	43	41	39
Wind Speed(kmph)	17	16	15	10	11
Wind Direction(Degree)	265	280	283	269	291
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Rainfall is expected for 1 day in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (61.0% & 41.0%), 36.0°C and 25.0°C respectively. The clear sky may appear. Expected average wind speed is 13.8 kmph. The Wind is expected to flow in west direction.

General Advisory:

Tamilnadu's average rainfall is likely to be increase than the normal during this week

SMS Advisory:

Blackgram VBN 8 & VBN 11 are available in KVK, Sirugamani, Trichy – 115. Contact: 8508835287

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing weather thrips infestation may occur in paddy at nursery stage. To control thrips infestation spray thiamethoxam 25% WG 4g
SUGARCANE	To control mealy bug in sugarcane, spray malathion 2ml/litre.
COTTON	Due to prevailing weather the white fly infestation may occur in cotton. To control white fly in cotton, Spray profenophos 50% EC 1000ml/ha.
MAIZE	Due to prevailing weather the downy mildew incidence may occur in maize. To control downy mildew, Spray mancozeb 1 kg/ha at 20 days after sowing.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Due to prevailing weather the white fly incidence may occur in Tomato. To control install yellow sticky traps @12/ha.
BANANA	Light to heavy wind/gust speed will expected in forthcoming days so farmers are advised to save the banana crop by using stacking method with casuarina poles.

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	Due to prevailing weather the Thrips incidence may occur in Chillies. To control, Spray Imidacloprid 17.8 SL 3 ml/10 litre.
JASMINE	For red spider mite incidence in jasmine, spraying of fenazagnin 10%EC @ 2ml/litre is recommended.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Pneumonia in goats – Fever, Respiratory distress, Mucous discharge from nostril, Reduced feed intake & Weight gain and Cough. Remedial measures: Provide clean water and ventilated house.
COW	To avoid mastitis disease in milching animals use potassium permanganate for cleaning the udder before and after milking.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Due to high wind speed feed material of poultries may be drifted hence hanging of gunny bags at the borders of the shed to reduce the wind speed.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 04-08-2023

Weather Forecast of **THOTTIAM** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2023-08-04(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-05	2023-08-06	2023-08-07	2023-08-08	2023-08-09
Rainfall	0.0	0.0	0.0	3.0	1.0
Tmax(°C)	36.0	36.0	34.0	36.0	37.0
Tmin(°C)	24.0	24.0	24.0	24.0	24.0
RH-I(%)	61	60	64	64	62
RH-II(%)	48	43	44	45	44
Wind Speed(kmph)	16	15	15	11	10
Wind Direction(Degree)	261	273	276	266	281
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Rainfall is expected for 2 day in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (62.0% & 45.0%), 36.0°C and 24.0°C respectively. The clear sky may appear. Expected average wind speed is 13.4 kmph. The Wind is expected to flow in west direction.

General Advisory:

Tamilnadu's average rainfall is likely to be increase than the normal during this week

SMS Advisory:

Blackgram VBN 8 & VBN 11 are available in KVK, Sirugamani, Trichy – 115. Contact: 8508835287

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing weather thrips infestation may occur in paddy at nursery stage. To control thrips infestation spray thiamethoxam 25% WG 4g
SUGARCANE	To control mealy bug in sugarcane, spray malathion 2ml/litre.
COTTON	Due to prevailing weather the white fly infestation may occur in cotton. To control white fly in cotton, Spray profenophos 50% EC 1000ml/ha.
MAIZE	Due to prevailing weather the downy mildew incidence may occur in maize. To control downy mildew, Spray mancozeb 1 kg/ha at 20 days after sowing.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Due to prevailing weather the white fly incidence may occur in Tomato. To control install yellow sticky traps @12/ha.
BANANA	Light to heavy wind/gust speed will expected in forthcoming days so farmers are advised to save the banana crop by using stacking method with casuarina poles.

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	Due to prevailing weather the Thrips incidence may occur in Chillies. To control, Spray Imidacloprid 17.8 SL 3 ml/10 litre.
JASMINE	For red spider mite incidence in jasmine, spraying of fenazagnin 10%EC @ 2ml/litre is recommended.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Pneumonia in goats – Fever, Respiratory distress, Mucous discharge from nostril, Reduced feed intake & Weight gain and Cough. Remedial measures: Provide clean water and ventilated house.
COW	To avoid mastitis disease in milching animals use potassium permanganate for cleaning the udder before and after milking.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Due to high wind speed feed material of poultries may be drifted hence hanging of gunny bags at the borders of the shed to reduce the wind speed.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 04-08-2023

Weather Forecast of **TURAIYUR** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2023-08-04(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-05	2023-08-06	2023-08-07	2023-08-08	2023-08-09
Rainfall	0.0	0.0	0.0	4.0	2.0
Tmax(°C)	36.0	35.0	34.0	35.0	36.0
Tmin(°C)	25.0	25.0	25.0	24.0	25.0
RH-I(%)	64	61	63	66	65
RH-II(%)	47	42	44	46	42
Wind Speed(kmph)	15	12	12	11	9
Wind Direction(Degree)	258	272	272	253	279
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Rainfall is expected for 2 day in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (64.0% & 44.0%), 35.0°C and 25.0°C respectively. The clear sky may appear. Expected average wind speed is 11.8 kmph. The Wind is expected to flow in west direction.

General Advisory:

Tamilnadu's average rainfall is likely to be increase than the normal during this week

SMS Advisory:

Blackgram VBN 8 & VBN 11 are available in KVK, Sirugamani, Trichy – 115. Contact: 8508835287

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing weather thrips infestation may occur in paddy at nursery stage. To control thrips infestation spray thiamethoxam 25% WG 4g
SUGARCANE	To control mealy bug in sugarcane, spray malathion 2ml/litre.
COTTON	Due to prevailing weather the white fly infestation may occur in cotton. To control white fly in cotton, Spray profenophos 50% EC 1000ml/ha.
MAIZE	Due to prevailing weather the downy mildew incidence may occur in maize. To control downy mildew, Spray mancozeb 1 kg/ha at 20 days after sowing.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Due to prevailing weather the white fly incidence may occur in Tomato. To control install yellow sticky traps @12/ha.
BANANA	Light to heavy wind/gust speed will expected in forthcoming days so farmers are advised to save the banana crop by using stacking method with casuarina poles.

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	Due to prevailing weather the Thrips incidence may occur in Chillies. To control, Spray Imidacloprid 17.8 SL 3 ml/10 litre.
JASMINE	For red spider mite incidence in jasmine, spraying of fenazagnin 10%EC @ 2ml/litre is recommended.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Pneumonia in goats – Fever, Respiratory distress, Mucous discharge from nostril, Reduced feed intake & Weight gain and Cough. Remedial measures: Provide clean water and ventilated house.
COW	To avoid mastitis disease in milching animals use potassium permanganate for cleaning the udder before and after milking.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Due to high wind speed feed material of poultries may be drifted hence hanging of gunny bags at the borders of the shed to reduce the wind speed.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 04-08-2023

Weather Forecast of **UPPILIAPURAM** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2023-08-04(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-05	2023-08-06	2023-08-07	2023-08-08	2023-08-09
Rainfall	0.0	0.0	0.0	5.0	2.0
Tmax(°C)	36.0	36.0	34.0	35.0	37.0
Tmin(°C)	24.0	24.0	25.0	24.0	24.0
RH-I(%)	64	62	65	67	66
RH-II(%)	48	43	45	47	44
Wind Speed(kmph)	15	12	12	10	8
Wind Direction(Degree)	258	271	272	255	278
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Rainfall is expected for 2 day in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (65.0% & 45.0%), 36.0°C and 24.0°C respectively. The clear sky may appear. Expected average wind speed is 11.4 kmph. The Wind is expected to flow in west direction.

General Advisory:

Tamilnadu's average rainfall is likely to be increase than the normal during this week

SMS Advisory:

Blackgram VBN 8 & VBN 11 are available in KVK, Sirugamani, Trichy – 115. Contact: 8508835287

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing weather thrips infestation may occur in paddy at nursery stage. To control thrips infestation spray thiamethoxam 25% WG 4g
SUGARCANE	To control mealy bug in sugarcane, spray malathion 2ml/litre.
COTTON	Due to prevailing weather the white fly infestation may occur in cotton. To control white fly in cotton, Spray profenophos 50% EC 1000ml/ha.
MAIZE	Due to prevailing weather the downy mildew incidence may occur in maize. To control downy mildew, Spray mancozeb 1 kg/ha at 20 days after sowing.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Due to prevailing weather the white fly incidence may occur in Tomato. To control install yellow sticky traps @12/ha.
BANANA	Light to heavy wind/gust speed will expected in forthcoming days so farmers are advised to save the banana crop by using stacking method with casuarina poles.

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	Due to prevailing weather the Thrips incidence may occur in Chillies. To control, Spray Imidacloprid 17.8 SL 3 ml/10 litre.
JASMINE	For red spider mite incidence in jasmine, spraying of fenazagnin 10%EC @ 2ml/litre is recommended.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Pneumonia in goats – Fever, Respiratory distress, Mucous discharge from nostril, Reduced feed intake & Weight gain and Cough. Remedial measures: Provide clean water and ventilated house.
COW	To avoid mastitis disease in milching animals use potassium permanganate for cleaning the udder before and after milking.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Due to high wind speed feed material of poultries may be drifted hence hanging of gunny bags at the borders of the shed to reduce the wind speed.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 04-08-2023

Weather Forecast of **VAIYAMPATTI** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2023-08-04(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-05	2023-08-06	2023-08-07	2023-08-08	2023-08-09
Rainfall	0.0	0.0	0.0	2.0	1.0
Tmax(°C)	35.0	35.0	33.0	34.0	35.0
Tmin(°C)	24.0	24.0	24.0	24.0	24.0
RH-I(%)	60	59	61	60	59
RH-II(%)	48	45	45	45	44
Wind Speed(kmph)	14	13	13	9	10
Wind Direction(Degree)	257	278	286	279	287
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Rainfall is expected for 2 day in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (60.0% & 45.0%), 34.0°C and 24.0°C respectively. The clear sky may appear. Expected average wind speed is 11.8 kmph. The Wind is expected to flow in west direction.

General Advisory:

Tamilnadu's average rainfall is likely to be increase than the normal during this week

SMS Advisory:

Blackgram VBN 8 & VBN 11 are available in KVK, Sirugamani, Trichy – 115. Contact: 8508835287

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing weather thrips infestation may occur in paddy at nursery stage. To control thrips infestation spray thiamethoxam 25% WG 4g
SUGARCANE	To control mealy bug in sugarcane, spray malathion 2ml/litre.
COTTON	Due to prevailing weather the white fly infestation may occur in cotton. To control white fly in cotton, Spray profenophos 50% EC 1000ml/ha.
MAIZE	Due to prevailing weather the downy mildew incidence may occur in maize. To control downy mildew, Spray mancozeb 1 kg/ha at 20 days after sowing.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Due to prevailing weather the white fly incidence may occur in Tomato. To control install yellow sticky traps @12/ha.
BANANA	Light to heavy wind/gust speed will expected in forthcoming days so farmers are advised to save the banana crop by using stacking method with casuarina poles.

Horticulture(Varieties)	Horticulture Specific Advisory
CHILLI	Due to prevailing weather the Thrips incidence may occur in Chillies. To control, Spray Imidacloprid 17.8 SL 3 ml/10 litre.
JASMINE	For red spider mite incidence in jasmine, spraying of fenazagnin 10%EC @ 2ml/litre is recommended.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Pneumonia in goats – Fever, Respiratory distress, Mucous discharge from nostril, Reduced feed intake & Weight gain and Cough. Remedial measures: Provide clean water and ventilated house.
COW	To avoid mastitis disease in milching animals use potassium permanganate for cleaning the udder before and after milking.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Due to high wind speed feed material of poultries may be drifted hence hanging of gunny bags at the borders of the shed to reduce the wind speed.