



Gramin Krishi Mausam Sewa
District Level Agromet Advisory Bulletin
 Tamil Nadu Agricultural University,
 Coimbatore, Tamil Nadu



Agromet Advisory Bulletin

Date : 03-10-2023

Weather Forecast of District RAMANATHAPURAM(Tamil Nadu) Issued On : 2023-10-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-10-04	2023-10-05	2023-10-06	2023-10-07	2023-10-08
Rainfall(mm)	0.0	0.0	0.0	0.0	3.0
Tmax(°C)	33.0	34.0	34.0	34.0	33.0
Tmin(°C)	25.0	24.0	24.0	24.0	24.0
RH-I(%)	70	70	70	70	70
RH-II(%)	50	50	50	50	50
Wind Speed(kmph)	22	20	18	16	18
Wind Direction(Degree)	200	200	200	180	180
Cloud Cover(Octa)	7	6	6	7	6

Weather Summary/Alert:

Light rainfall expected for 8th October. Maximum temperature would be around 33-34°C and the minimum temperature would be around 24-25°C. Relative humidity can expected the range of 50-70 percent. Partly cloudy sky will appear for the next five days. Wind speed of 16-22 kmph expected from South directions

General Advisory:

From extended range forecast, above normal rainfall, normal maximum and minimum temperature expected for the period of October 08 to October 14 over Tamil Nadu.

SMS Advisory:

Farmers are advised to apply 200 kg of gypsum per acre at the time of last ploughing for better growth of crop.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	In rainfed rice cultivation, it is recommended for farmers to apply 100kg/ac of single super phosphate as a basal dose.
RICE	In case of rainfed direct seeding of paddy, the use of mechanized sowing facilitate the possibility of maintaining the proper density spacing and minimize the crop loss due to dry spell. One kg of seed should be soaked in a one per cent potassium permanganate solution (10g in a liter of water) for 16 hours and then dried in the shade until it returns to its original moisture content.

Horticulture Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE RICE	phosphate as a basal dose. In case of rain fed direct seeding of paddy, the use of mechanized sowing facilitate the possibility of maintaining the proper depth spacing and minimize the crop loss due to dry hardening. one kg of seed should be soaked in a one per cent potassium permanganate solution (10 g in a liter of water) for 16 hours and then dried in the shade until it returns to its original moisture content.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	Due to the prevailing weather condition, incidence of red palm weevil expected in coconut. The damage symptoms are holes can be seen on the stem with chewed up fibres protruding out and reddish brown liquid can be seen oozing out from the hole. If symptoms are observed in coconut tree, spot application with insecticidal solution containing Imidacloprid 17.8% SL @ 1 ml with 1 liter of water.
COCONUT	In coconut, to control the Rhinocerosbeetle, the topmost three leaf axils around the spindle filled with Naphthalene balls three to four balls covered with fine sand, once in 45 days. Soak castor cake at 1 kg in 5 liter of water in small mud pots and keep them in the coconut gardens to attract and kill the adults.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	As per the advice of the veterinary doctor, farmers are instructed to vaccinate their dairy animals against Haemorrhagic septicaemia (HS), Black Quarter (BQ) and FMD disease