



Agromet Advisory Bulletin for the **Nagaipattinam** District

(From 19th April 2024 to 23rd April 2024)

Issued jointly by Tamil Nadu Rice Research Institute -
Aduthurai and IMD



Bulletin no: 06/2024

Date: 18.04.2024

Weather Summary for the period since the issue of last bulletin no. 05 (16.04.2024) for Tamil Nadu: Nagaipattinam district

Maximum temperature of 34 - 35 °C, minimum temperature of 26 - 27 °C, was recorded in Nagaipattinam district.

Weather forecast for next five days (19.04.2024 to 23.04.2024)

Past week Summary 16.04.2024 to 17.04.2024		Parameters	Ensemble weather forecast valid until 08.30 hr of 23.04.2024				
Day-1 16/04	Day-2 17/04		Day-1 19/04	Day-2 20/04	Day-3 21/04	Day-4 22/04	Day-5 23/04
0	0	Rainfall (mm)	0	0	0	1	1
34	35	Max. Temp. (°C)	35	36	36	37	36
27	26	Min. Temp. (°C)	26	27	26	27	26
5	4	Cloud cover (Octa)	3	1	3	5	6
75-74	80-78	RH I-II (%)	80-50	80-50	80-50	80-50	70-50
6	5	Wind (kmph)	22	22	20	20	14
North East	North East	Wind direction	South	South	South	South	South South East

Weather Summary and forecast for Nagaipattinam district

According to district level forecast issued by India Meteorological Department for Nagaipattinam district, the sky will be clear. Light rainfall expected on 22.04.2024 and 23.04.2024. Maximum temperature is expected to be around 35 - 37 °C and minimum temperature is expected to be around 26 - 27 °C. Morning relative humidity is expected to be around 70 - 80 percent and evening relative humidity is expected to be around 50 percent. Average wind speed is expected to be around 14 - 22 km per hour. The wind direction will be south.

Agro Advisory

Crop	Stage	Advisory
------	-------	----------

General		The prevailing weather condition favors for summer ploughing.
Pulses	Harvest	Harvest the matured blackgram and greengram plants.
		Harvested plants of blackgram and greengram plants can be sundried before storage.
Coconut		Summer, root feeding of coconut tonic @ 200 ml/palm should be done. To control & decreases the button shedding of coconut.
Cattle	Cow	During summer, tick infestation flares up in all livestock. This might cause anemia & makes up way for various disease occurrences. The animal will be dull & depressed, difficulty in feed intake. Mix 3ml of butox in 1 litre of water and wash the animals with damp cloth.

SMS advisory

The prevailing weather condition favors for summer ploughing.

**Director
TRRI, Aduthurai**