

Agromet Advisory Bulletin for the Nagaipattinam District

(From 04th September 2024 to 08th September 2024)



Issued jointly by Tamil Nadu Rice Research Institute - Aduthurai and IMD

Bulletin no: 45/2024

Date: 03.09.2024

Weather Summary for the period since the issue of last bulletin no. 44 (30.08.2024) for Tamil Nadu: Nagaipattinam district

Maximum temperature of 35 - 37 $^{\circ}$ C, minimum temperature of 23 - 24 $^{\circ}$ C, and isolated rainfall was recorded in Nagai pattinam district.

Weather forecast for next five days (04.09.2024 to 08.09.2024)

Past week Summary 30.08.2024 to 02.09.2024				Parameters	Ensemble weather fore cast valid until 08.30 hr of 08.09.2024				
Day-1	Day-2	Day-3	Day-4		Day-1	Day-2	Day-3	Day-4	Day-5
30/08	31/08	01/09	02/09	Date	04/09	05/09	06/09	07/09	08/09
13.4	0.0	1.6	0.0	Rainfall (mm)	2	3	2	0	8
35	35	37	37	Max. Temp. (°C)	38	38	37	37	36
24	24	24	23	Min. Temp. (°C)	27	27	26	27	26
7	8	6	6	Cloud cover (Octa)	8	7	6	6	7
79-71	74-68	64-61	65-49	RH I-II (%)	60-40	60-40	70-40	70-40	70-40
7	7	11	7	Wind (kmph)	18	20	22	20	22
South	South South West	South	South	Wind direction	South West	South West	South South West	South South West	South South West

Weather Summary and forecast for Nagaipattinam district

According to district level forecast issued by India Meteorological Department for Nagaipattinam district, the sky will be clear. Light drizzling expected on 04.09.2024, 05.09.2024, 06.08.2024 and 08.09.2024. Maximum temperature is expected to be around 36 - 38 °C and minimum temperature is expected to be around 26 - 27 °C. Morning relative humidity is expected to be around 40 percent. Average wind speed is expected to be around 18 - 22 km per hour. The wind direction will be south south west.

Agro Advisory

Crop	Stage	Advisory

		Farmers are advised to provide adequate drainage facilities to all		
		crops due to light rainfall forecast.		
Gei	neral	Farmers are requested to download and utilize the agricultural		
		based mobile apps suggested by TNAU (TNAU AAS, Meghdoot and		
		DAMINI).		
	Nursery	For rice nursery, applications of 2 kg DAP/cent is recommended. In		
Rice		rice nursery, spray Azadiraction 0.03% (3 ml/litre of water) with		
	riarsery	sticking agent Khadi Gopura Bar Soap @ 3g/litre to prevent thrips		
		infection on 10 days after sowing.		
	Cow	This situation is favorable for the development of Aflatoxin		
Cattle		infection in the Corn kernels and Paddy Straw, which may affect		
Gattle		the health of the cattle. To save them from infection, dry the Corn		
		kernels and Paddy Straw before feeding.		

SMS advisory

Farmers are advised to provide adequate drainage facilities to all crops due to light rainfall forecast.

Director TRRI, Aduthurai